Optimization and Control Under Uncertainty

Drew Kouri

Optimization and Uncertainty Quantification
Sandia National Laboratories, Albuquerque, New Mexico
dpkouri@sandia.gov

April 19, 2018

SIAM Conference on Uncertainty Quantification
Garden Grove, CA.

Funding: DARPA EQUiPS

Sandia National Laboratories is a multimission laboratory managed and operated by National Technology and Engineering Solutions of Sandia, LLC., a wholly owned subsidiary of Honeywell International, Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA0003525.

Outline

Motivating Applications

General Problem Formulation

Quantifying Risk

Probabilistic Optimization

What if our uncertainty is uncertain?

Computational Solution Methods

Software

References

Topology Optimization & Additive Manufacturing



Given $V_0 \in (0,1)$ compute a density that solves:

$$\underset{0 \le z \le 1}{\text{Minimize}} \; \mathcal{R}\left(\int_D F \cdot S(z)\, dx + \int_{\Gamma_t} t \cdot S(z)\, dx \right)$$

s.t. $\int_D z(x)\, dx \le V_0 |D|$, where $S(z) = u$ solves the **linear elasticity equations**

$$\begin{aligned} -\nabla \cdot (\mathbf{E}(z) : \epsilon u) &= F, && \text{in } D, \text{ a.s.} \\ \epsilon u &= \frac{1}{2}(\nabla u + \nabla u^\top), && \text{in } D, \text{ a.s.} \\ \epsilon u n &= t, && \text{on } \Gamma_t, \text{ a.s.} \\ u &= g, && \text{on } \Gamma_d, \text{ a.s.} \end{aligned}$$

- Uncertain external forces (loads) and boundary conditions.
- Uncertain internal forces, e.g., residual stresses due to AM.
- Uncertain material properties (porosity, etc.) due to AM.
- **Reliability formulation:** Compute light-weight designs that minimize the probability of structural failure.

Reservoir Optimization: Secondary Oil Recovery



Given $D \subset \mathbb{R}^3$ and interest rate $r \geq 0$:

$$\underset{z\,=\,(q,\,\widehat{q})}{\text{Minimize}}\ \mathcal{R}\left(\int_0^T e^{rt}\, C([S(z)](t), z(t), t)\, dt\right)$$

where $S(z) = (s, v, p)$ solves the **reservoir equations**

$$\begin{aligned} -\mathbf{K}\lambda(s)\nabla p &= v, && \text{in } D,\ \text{a.s.} \\ \nabla \cdot v &= q, && \text{in } D,\ \text{a.s.} \\ \phi\, \partial_t s + \nabla \cdot (f(s) v\,) &= \widehat{q}, && \text{in } D,\ \text{a.s.} \end{aligned}$$

(plus initial and boundary conditions).

- Porosity, ϕ, and permeability, $\mathbf{K}$, are estimated from data (e.g., seismic inversion).
- Total mobility, λ, and fractional flow function, f, may be uncertain.
- **Risk-neutral formulation:** Determine injection rates that minimize cost on average.
- **Risk-averse formulation:** Determine injection rates that minimize the average of the 10% worst costs.

Control of Chemical Vapor Deposition Reactors





Consider the optimal control problem

$$\min_z \ \frac{1}{2}\mathcal{R}\left(\int_D (\nabla \times U(z))\,\mathrm{d}x\right) + \frac{\gamma}{2}\int_{\Gamma_c} |z|^2\,\mathrm{d}x$$

where $S(z) = (U(z), P(z), T(z)) = (u, p, t)$ solves the **Boussinesq flow equations**

$$\begin{aligned} -\nu\nabla^2 u + (u\cdot\nabla)u + \nabla p + \eta t g &= 0 \quad \text{in } D, \text{ a.s.} \\ \nabla\cdot u &= 0 \quad \text{in } D, \text{ a.s.} \\ -\kappa\Delta t + u\cdot\nabla t &= 0 \quad \text{in } D, \text{ a.s.} \\ \kappa\nabla t\cdot n + h(z-t) &= 0 \quad \text{on } \Gamma_c, \text{ a.s.} \end{aligned}$$

(plus additional boundary conditions).

- Uncertain viscosity, thermal conductivity, substrate temperature, etc. imply flow velocity, pressure and temperature are uncertain.
- **Risk-averse formulation:** Determine wall temperature that minimizes the average of *low-probability*, large vorticity scenarios.

Direct Field Acoustic Testing



Consider the optimal control problem

$$\min_z \ \frac{1}{2}\mathcal{R}\left(\int_{D_o} (U(z)-w)\overline{(U(z)-w)}\,\mathrm{d}x\right)+\frac{\gamma}{2}\int_{D_c} |z|^2\,\mathrm{d}x$$

where $U(z) = u$ solves the **Helmholtz equation**

$$\begin{aligned} -\Delta u - \kappa^2(1+\sigma\epsilon)^2 u &= \mathbb{1}_{D_c} z \qquad \text{in } D, \text{ a.s.} \\ \nabla u \cdot n &= i\kappa u \qquad \text{on } \partial D, \text{ a.s.} \end{aligned}$$

- The refractive index of the device under investigation is often uncertain.
- **Risk-neutral formulation:** Determine speaker output that produces a material response that matches a desired vibration profile on average.
- **Risk-averse formulation:** Determine speaker output that produces a response that is "good" on average for the 10% worst scenarios.

Optimizing Your Short Game

Vanderbei 2001

Determine initial velocity yeilding a trajectory with minimal speed at the hole:

$$\min_z \ \mathcal{R}\left(\frac{1}{2}(v_1(T)^2 + v_2(T)^2)\right)$$

where $S(z) = (x, v, a, T)$ solves the equations of motion

$$ma = N + F - mge_3$$
$$\dot{x} = v, \quad \dot{v} = a$$
$$x(0) = x, \quad v(0) = Bz$$

where the normal $N = N(x, v, a)$ and friction $F = F(x, v, a)$ forces are

$$N_i = -\frac{\partial x_3}{\partial x_i} N_3, \quad N_3 = m\frac{g - a_1\frac{\partial x_3}{\partial x_1} - a_2\frac{\partial x_3}{\partial x_2} + a_3}{(\partial x_3/\partial x_1)^2 + (\partial x_3/\partial x_2)^2 + 1} \quad \text{and} \quad F = -\mu\|N\|\frac{v}{\|v\|}$$



Outline

Motivating Applications

General Problem Formulation

Quantifying Risk

Probabilistic Optimization

What if our uncertainty is uncertain?

Computational Solution Methods

Software

References

General PDE-Optimization under Uncertainty

Making Deterministic Problems Stochastic

Deterministic PDE-Constrained Optimization:

U and Z are reflexive Banach spaces, Z_{ad} is a closed convex subset of Z, Y is a Banach space, $J : U \times Z \to \mathbb{R}$ and $c : U \times Z \to Y$:

$$\underset{z \in Z_{\text{ad}}}{\text{Minimize}} \ \widehat{J}(z)$$

where $\widehat{J}(z) := J(S(z), z)$ and $S(z) = u \in U$ solves the PDE

$$c(u, z) = 0.$$

Stochastic PDE-Constrained Optimization:

$(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space. Objective function and PDE are now **parametrized**, i.e., $J : U \times Z \times \Omega \to \mathbb{R}$ and $c : U \times Z \times \Omega \to Y$:

$$\underset{z \in Z_{\text{ad}}}{\text{Minimize}} \ \mathcal{J}(z) = \mathcal{R}(\widehat{J}(z))$$

where $\widehat{J}(z) := J(S(z), z, \cdot)$ and $S(z) = u : \Omega \to U$ solves the PDE

$$c(u, z, \omega) = 0.$$

Notation

$(\Omega, \mathcal{F})$ is a measurable space

$\mathbb{P}$, $P : \mathcal{F} \to [0,1]$ are probability measures

1. **Expectation:** $\mathbb{E}_P[X] = \int_\Omega X(\omega)\,\mathrm{d}P(\omega)$ and $\mathbb{E}[X] = \mathbb{E}_\mathbb{P}[X]$
2. **Variance:** $\mathbb{V}_P[X] = \mathbb{E}_P[(X - \mathbb{E}_P[X])^2]$ and $\mathbb{V}[X] = \mathbb{V}_\mathbb{P}[X]$
3. **Standard Deviation:** $\sigma_P[X] = \mathbb{V}_P[X]^{1/2}$ and $\sigma[X] = \sigma_\mathbb{P}[X]$
4. **Distribution:** $F_X(x) = \mathbb{P}(X \leq x)$
5. **Quantile:** $q_\beta(X) = \inf\{t \in \mathbb{R} \mid F_X(x) > \beta\} = F_X^{-1}(\beta)$

Tensor Product Function Spaces

Lebesgue Spaces: For $1 \leq p < \infty$,

$$L^p(\Omega, \mathcal{F}, \mathbb{P}) := \left\{ v : \Omega \to \mathbb{R} \;\mid\; v\ \mathcal{F}\text{-measurable}, \int_\Omega |v(\omega)|^p \, \mathrm{d}\mathbb{P}(\omega) < \infty \right\},$$

$$L^\infty(\Omega, \mathcal{F}, \mathbb{P}) := \{ v : \Omega \to \mathbb{R} \;\mid\; v\ \mathcal{F}\text{-measurable}, \operatorname{ess\,sup} |v(\omega)| < \infty \}.$$

If $f,\, g \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ then $f = g \iff f(\omega) = g(\omega)$ for $\mathbb{P}$ almost all $\omega \in \Omega$.

Tensor Spaces: Given a real Banach space W then

$$L^p(\Omega, \mathcal{F}, \mathbb{P}) \otimes W := \text{span}\, \{ vx \;\mid\; v \in L^p(\Omega, \mathcal{F}, \mathbb{P}),\ x \in W \}.$$

Many norms exist for the vector space $L^p(\Omega, \mathcal{F}, \mathbb{P}) \otimes W$ and given a norm $L^p(\Omega, \mathcal{F}, \mathbb{P}) \otimes W$ is not necessarily complete.

Bochner Spaces: For $1 \leq p < \infty$ and W a real Banach space

$$L^p(\Omega, \mathcal{F}, \mathbb{P}; W) := \left\{ v : \Omega \to W \;\mid\; v \text{ strongly } \mathcal{F}\text{-measurable}, \int_\Omega \|v(\omega)\|_W^p \, \mathrm{d}\mathbb{P}(\omega) < \infty \right\}$$

and similarly for $p = \infty$. $L^p(\Omega, \mathcal{F}, \mathbb{P}; W)$ is the completion of $L^p(\Omega, \mathcal{F}, \mathbb{P}) \otimes W$ with respect to the Bochner norm

$$\|u\|_{L^p(\Omega,\mathcal{F},\mathbb{P};W)} := \left(\int_\Omega \|u(\omega)\|_W^p \, \mathrm{d}\mathbb{P}(\omega) \right)^{\frac{1}{p}} \quad \text{and} \quad \|u\|_{L^\infty(\Omega,\mathcal{F},\mathbb{P};W)} := \operatorname{ess\,sup} \|u(\omega)\|_W.$$

Again, if $f,\, g \in L^p(\Omega, \mathcal{F}, \mathbb{P}; W)$ then $f = g \iff f(\omega) = g(\omega)$ for $\mathbb{P}$ almost all $\omega \in \Omega$.

Assumptions on PDE Solution Map $S(z)$

1. For each $z \in Z$, $c(u, z, \omega) = 0$ is well posed, i.e.,
   (i) $\exists!\, S(z) : \Omega \to U$ such that $c(S(z), z, \cdot) = 0$ a.s. for all z;
   (ii) $\exists\, 0 < c(\cdot) \in L^q(\Omega, \mathcal{F}, \mathbb{P})$, $1 \leq q \leq \infty$ and an increasing function $\rho : [0, \infty) \to [0, \infty)$ both independent of z such that

$$\|S(z)\|_U \leq c\rho(\|z\|_Z) \quad \text{a.s.} \quad \forall z \in Z_{\text{ad}}.$$

2. $S(z)$ is strongly measurable $\forall z \in Z_{\text{ad}} \implies S(z) \in L^q(\Omega, \mathcal{F}, \mathbb{P}; U)$.
3. $z \mapsto S(z)$ satisfies the continuity property

$$z_n \rightharpoonup z \ \text{ in } \ Z \quad \implies \quad S(z_n) \rightharpoonup S(z) \ \text{ in } \ U, \ \text{a.s.}$$

4. $\exists V \supseteq Z_{\text{ad}}$, V open, such that $S : V \to L^q(\Omega, \mathcal{F}, \mathbb{P}; U)$ is continuously Fréchet differentiable.

Senstivity Equation: To compute the sensitivity of $S(z)$ in the direction $h \in Z$ solve:

$$c_u(S(z), z, \cdot)S'(z)h + c_z(S(z), z, \cdot)h = 0 \quad \text{a.s.}$$

Example: Linear Elliptic PDE

Let $D \subset \mathbb{R}^n$ be a bounded Lipschitz domain, $U = H_0^1(D)$, $Y = Z = H^{-1}(D)$ and $A : \Omega \to \mathbb{R}^{n \times n}$:

$$\langle c(u, z, \omega), v \rangle_{U^*, U} := \int_D (A(\omega)\nabla u(x)) \cdot \nabla v(x)\, \mathrm{d}x - \langle z, v \rangle_{U^*, U} \quad \text{for } v \in H_0^1(D).$$

If $\exists\, 0 < \underline{c} \leq \bar{c} < \infty$ such that

$$\underline{c} \leq \frac{\zeta^\top A(\omega)\zeta}{\zeta^\top \zeta} \leq \bar{c} \quad \text{a.s.}$$

then Lax-Milgram $\implies$ existence of a unique solution $u \in H_0^1(D)$ to $c(u, z, \cdot) = 0$ for fixed z a.s. Moreover,

$$\underline{c}\|\nabla S(z)\|_{L^2(D)}^2 \leq \|z\|_{H^{-1}(D)} \|S(z)\|_{H_0^1(D)} \quad \text{a.s.}$$

Hence, Poincaré's inequality guarantees that

$$\|\nabla S(z)\|_{L^2(D)} \leq C_{d,D} \|z\|_{H^{-1}(D)} \quad \text{a.s.}$$

and $S : H^{-1}(D) \to L^\infty(\Omega, \Sigma, \mathbb{P}; H_0^1(D))$.

Note: S with domain restricted to $L^2(D)$ is compact since $L^2(D) \subset\subset H^{-1}(D)$.

Uncertain Objective Functions

General Assumptions:

1. **Integrability:** $\widehat{J}(z) \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ for all $z \in Z$;
2. **Weak Lower Semicontinuity:** If $z_n \rightharpoonup z$ then

$$\liminf_{n \to \infty} \mathbb{E}[\vartheta \widehat{J}(z_n)] \geq \mathbb{E}[\vartheta \widehat{J}(z)]$$

for all $\vartheta \in (L^p(\Omega, \mathcal{F}, \mathbb{P}))^*$ satisfying $\vartheta \geq 0$ a.s.

Compare to *normal integrands*, i.e., the epigraph of $\widehat{J}$ is measurable and closed valued.

Uncertain Objective Functions

Separable Objective Functions: $J(u, z, \omega) = g(u, \omega) + \wp(z)$

1. **Carathéodory:** $g(\cdot, \omega)$ is continuous a.s. and $g(u, \cdot)$ is measurable $\forall\, u \in U$.
2. **Growth Condition:**
   If $q < \infty$, then $\exists\, 0 \leq a \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ and $c > 0$ such that

   $$|g(u, \omega)| \leq a(\omega) + c\|u\|_U^{q/p} \quad \forall\, u \in U \text{ a.s.}$$

   If $q = \infty$, then $\forall\, c > 0 \;\; \exists\, \gamma_c \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ such that

   $$|J(u, \omega)| \leq \gamma_c(\omega) \quad \text{a.s.} \quad \forall\, u \in U, \;\; \|u\|_U \leq c.$$

3. **Convexity:** $g(\cdot, \omega)$ is convex a.s. (optional)

Uncertain Objective Functions

The Separable Case

Superposition (Nemytskii) Operator:
$\mathcal{G} : L^q(\Omega, \mathcal{F}, \mathbb{P}; U) \to L^p(\Omega, \mathcal{F}, \mathbb{P})$ where $\mathcal{G}(u) = g(u(\cdot), \cdot)$.

1. If g is Carathéodory and satisfies the growth condition, then $\mathcal{G} : L^q(\Omega, \mathcal{F}, \mathbb{P}; U) \to L^p(\Omega, \mathcal{F}, \mathbb{P})$ is continuous.
2. If, in addition, g is convex, then $\mathcal{G}$ is Gâteaux directionally differentiable.
3. If, in addition, g is locally Lipschitz, then $\mathcal{G}$ is Hadamard directionally differentiable.
4. If $g(\cdot, \omega)$ is continuously Fréchet differentiable for a.s. and there exists $\alpha > 0$ and $K \in L^s(\Omega, \mathcal{F}, \mathbb{P})$ with

$$s = \begin{cases} pq/(q - (1+\alpha)p) & \text{if } q > (1+\alpha)p \\ \infty & \text{if } q = (1+\alpha)p \end{cases}$$

such that

$$\|g_u(u, \omega) - g_u(v, \omega)\|_{U^*} \leq K(\omega)\|u - v\|_U^\alpha \quad \text{a.s.}$$

Then $\mathcal{G}$ is Fréchet differentiable.

Example: Quadratic Objective Function

Let W be a real Hilbert space, $w \in W$ and $\mathbf{C} \in \mathcal{L}(U, W)$. Consider

$$J(u, z, \omega) = \frac{1}{2}\|\mathbf{C}u - w\|_W^2 + \frac{\gamma}{2}\|z\|_Z^2, \quad \gamma > 0.$$

J is separable with $g(u, \omega) = \frac{1}{2}\|\mathbf{C}u - w\|_W^2$.

1. **Carathéodory:** Satisfied since g has no dependence on ω.
2. **Growth Condition:** Satisfied (using Young's inequality) with

$$a = \|w\|_W^2 \quad \text{and} \quad c = \|\mathbf{C}\|_{\mathcal{L}(U,W)}^2.$$

3. **Convexity:** Clearly satisifed.
4. **Differentiability:** Satisfied with $K = \|\mathbf{C}\|_{\mathcal{L}(U,W)}^2$ and $\alpha = 1$.

Result: $\mathcal{G} : L^q(\Omega, \mathcal{F}, \mathbb{P}; U) \to L^p(\Omega, \mathcal{F}, \mathbb{P})$ is continuous and Fréchet differentiable as long as $q \geq 2p$.

The Functional $\mathcal{R}$

Assumptions & Existence of Minimizers

$\mathcal{R} : L^p(\Omega, \mathcal{F}, \mathbb{P}) \to \mathbb{R} \cup \{+\infty\}$

- $\mathcal{R}$ is proper, convex and lower semicontinuous
- $\mathcal{R}$ satisfies $\mathcal{R}(C) = C$ for all constants C;
- $\mathcal{R}$ is monotonic, i.e., if $X \geq X'$ a.s., then $\mathcal{R}(X) \geq \mathcal{R}(X')$.

Existence: If Z_{ad} is convex, closed and bounded, then there exists a minimizer of $\mathcal{J}(z) = \mathcal{R}(\widehat{J}(z))$ in Z_{ad}.

Proof: Apply the direct method of the calculus of variations.

Note: The same result holds if $Z = Z_{\text{ad}}$ and $\widehat{J}(z)$ is a.s. coercive, i.e., $Z_{\text{ad}} = Z$ and $\widehat{J}(z)$ has the coercivity propery that $\exists\, r > 0$ and coercive $\varphi : Z \to \mathbb{R} \cup \{+\infty\}$, such that

$$\|z\|_Z \geq r \quad \Longrightarrow \quad \widehat{J}(z) \geq \varphi(z) \text{ a.s.}$$

The Functional $\mathcal{R}$

Assumptions & Existence of Minimizers

$\mathcal{R} : L^p(\Omega, \mathcal{F}, \mathbb{P}) \to \mathbb{R} \cancel{\cup \{+\infty\}}$

- $\mathcal{R}$ is ~~proper,~~ convex and lower semicontinuous
- $\mathcal{R}$ satisfies $\mathcal{R}(C) = C$ for all constants C;
- $\mathcal{R}$ is monotonic, i.e., if $X \geq X'$ a.s., then $\mathcal{R}(X) \geq \mathcal{R}(X')$.

Existence: If Z_{ad} is convex, closed and bounded, then there exists a minimizer of $\mathcal{J}(z) = \mathcal{R}(\widehat{J}(z))$ in Z_{ad}.

Proof: Apply the direct method of the calculus of variations.

Note: The same result holds if $Z = Z_{\text{ad}}$ and $\widehat{J}(z)$ is a.s. coercive, i.e., $Z_{\text{ad}} = Z$ and $\widehat{J}(z)$ has the coercivity propery that $\exists\, r > 0$ and coercive $\varphi : Z \to \mathbb{R} \cup \{+\infty\}$, such that

$$\|z\|_Z \geq r \quad \Longrightarrow \quad \widehat{J}(z) \geq \varphi(z) \text{ a.s.}$$

The Functional $\mathcal{R}$

Assumptions & Existence of Minimizers

$\mathcal{R} : L^p(\Omega, \mathcal{F}, \mathbb{P}) \to \mathbb{R} \cancel{\cup \{+\infty\}}$

- $\mathcal{R}$ is ~~proper,~~ convex and lower semicontinuous
- $\mathcal{R}$ satisfies $\mathcal{R}(C) = C$ for all constants C;
- $\mathcal{R}$ is monotonic, i.e., if $X \geq X'$ a.s., then $\mathcal{R}(X) \geq \mathcal{R}(X')$.

Existence: If Z_{ad} is convex, closed and bounded, then there exists a minimizer of $\mathcal{J}(z) = \mathcal{R}(\widehat{J}(z))$ in Z_{ad}.

Proof: Apply the direct method of the calculus of variations.

Note: The same result holds if $Z = Z_{\text{ad}}$ and $\widehat{J}(z)$ is a.s. coercive, i.e., $Z_{\text{ad}} = Z$ and $\widehat{J}(z)$ has the coercivity propery that $\exists\, r > 0$ and coercive $\varphi : Z \to \mathbb{R} \cup \{+\infty\}$, such that

$$\|z\|_Z \geq r \quad \Longrightarrow \quad \widehat{J}(z) \geq \varphi(z) \text{ a.s.}$$

Note: Need additional/different assumptions if $p = \infty$!

Outline

Motivating Applications

General Problem Formulation

Quantifying Risk

Probabilistic Optimization

What if our uncertainty is uncertain?

Computational Solution Methods

Software

References

Modeling Risk Preference

What is risk? *Possibility of loss or injury* (Merriam Webster)

. . . In our optimization problem, $J(S(z;\cdot),\cdot)$ is a **risk**!

We **cannot** directly minimize $J(S(z;\cdot),\cdot) + \wp(z) \in \mathcal{X} := L^p(\Omega, \mathcal{F}, \mathbb{P})$

. . . How should we quantify our risk?

- **Traditional Stochastic Programming:** Minimize *on average*

$$\mathcal{R}(\widehat{J}(z)) = \mathbb{E}[\widehat{J}(z)].$$

- **Risk-Averse Stochastic Programming:** Model *risk preferences*

$$\mathcal{R}(\widehat{J}(z)) = \mathbb{E}[\widehat{J}(z)] + c\mathbb{E}[(\widehat{J}(z) - \mathbb{E}[\widehat{J}(z)])_+^p]^{1/p}.$$

- **Probabilistic Optimization:** Minimize the *probability of loss*

$$\mathcal{R}(\widehat{J}(z)) = \mathbb{P}(\widehat{J}(z) > \tau).$$

- **Stochastic Orders:** Model risk preference with a *benchmark* Y

$$\mathbb{P}(\widehat{J}(z) \leq x) \leq \mathbb{P}(Y \leq x) \quad \forall\, x \in \mathbb{R}.$$

Quantifying Risk & Controlling Uncertainty

- Reduce **variability** of optimized system:

$$\mathbb{E}[(X - \mathbb{E}[X])^2] \qquad \text{or} \qquad \mathbb{E}[(X - \mathbb{E}[X])_+^p]^{1/p}$$

- Control **rare events**, reduce **failure**, and certify **reliability**:

$$\mathbb{P}(X > t) \qquad \text{or} \qquad q_\beta(X) = \inf\{\, t \in \mathbb{R} \,:\, \mathbb{P}(X \le t) \ge \beta \,\}$$

- Minimize over **undesirable events**:

$$\mathrm{CVaR}_\beta(X) = \frac{1}{1-\beta}\int_\beta^1 F_X^{-1}(\alpha)\,\mathrm{d}\alpha \approx \mathbb{E}[X \,|\, X \ge q_\beta(X)]$$



Mitigating Uncertainty by Shaping Distributions

Law Invariance & Stochastic Dominance

Law Invariance:

- $\mathcal{R}$ is **law invariant** if

$$F_X(t) = F_{X'}(t) \;\; \forall t \in \mathbb{R} \quad \Longrightarrow \quad \mathcal{R}(X) = \mathcal{R}(X').$$

  If $\mathcal{R}$ is law invariant, then it is a function of distributions.

Stochastic Dominance:

- X **dominates** X' with respect to the 1st **stochastic order**, denoted $X \succeq_{(1)} X'$, if

$$F_X(t) \leq F_{X'}(t) \;\; \forall t \in \mathbb{R}.$$

- X **dominates** X' with repsect to the 2nd **stochastic order**, denoted $X \succeq_{(2)} X'$, if

$$\int_{-\infty}^{t} F_X(\eta)\,\mathrm{d}\eta \leq \int_{-\infty}^{t} F_{X'}(\eta)\,\mathrm{d}\eta \quad \forall t \in \mathbb{R}$$
$$\Longleftrightarrow \quad \mathbb{E}[(t-X)_+] \leq \mathbb{E}[(t-X')_+] \;\; \forall t \in \mathbb{R}.$$

  Here, $(x)_+ = \max\{0, x\}$.

Consequences: Suppose $\mathcal{R}$ is law invariant:

- If $X \geq X'$ a.s. implies $\mathcal{R}(X) \geq \mathcal{R}(X')$, then $X \succeq_{(1)} X'$ implies $\mathcal{R}(X) \geq \mathcal{R}(X')$;
- If $\mathcal{R}$ is lsc and convex, then $-X' \succeq_{(2)} -X$ implies $\mathcal{R}(X) \geq \mathcal{R}(X')$.
- Law invariant $\mathcal{R}$ prefer dominated random variables!

Mean-Plus-Variance Risk

Markowitz, Portfolio Selection, 1952

A common risk functional in engineering application is

$$\mathcal{R}(X) = \mathbb{E}[X] + c\mathbb{V}[X] \quad \text{for } c > 0.$$

Downsides:

- $\mathcal{R}$ penalizes variation **below** the mean.
- $\mathcal{R}$ is **not** monotonic.

Example: Shapiro, Dentcheva, Ruszczynski (2014)

Suppose $\Omega = \{\omega_1, \omega_2\}$ with associated probabilities $\mathfrak{p} \in (0,1)$ and $(1-\mathfrak{p})$. Consider the stochastic program

$$\underset{z_1,\, z_2}{\text{Minimize}} \ \mathcal{R}(-\zeta_1 z_1 - \zeta_2 z_2) \quad \text{subject to} \quad z_1 + z_2 = 1 \ \text{ and } \ z_1,\, z_2 \geq 0$$

where $\zeta_1,\, \zeta_2 : \Omega \to \mathbb{R}$ are

$$\zeta_1(\omega_1) = a > 0, \quad \zeta_1(\omega_2) = 0, \quad \text{and} \quad \zeta_2(\omega_1) = \zeta_2(\omega_2) = 0.$$

If $\mathfrak{p} \leq 1 - (ca)^{-1}$, then $\mathcal{R}(-\zeta_1) = -\mathfrak{p}a + ca^2\mathfrak{p}(1-\mathfrak{p}) > \mathcal{R}(-\zeta_2) = 0$ eventhough $-\zeta_1 \leq -\zeta_2$ for all $\omega \in \Omega$.

Coherent Risk Measures

$\mathcal{R} : L^p(\Omega, \mathcal{F}, \mathbb{P}) \to \mathbb{R}$ is **coherent** if

(R1) **Subadditivity:** For all $X,\ X' \in L^p(\Omega, \mathcal{F}, \mathbb{P})$,

$$\mathcal{R}(X + X') \leq \mathcal{R}(X) + \mathcal{R}(X')$$

(R2) **Monotonicity:** For any $X,\ X' \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ satifying

$$X \geq X' \ \text{ a.s.} \qquad \Longrightarrow \qquad \mathcal{R}(X) \geq \mathcal{R}(X')$$

(R3) **Translation Equivariance:** For all $X \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ and $t \in \mathbb{R}$,

$$\mathcal{R}(X + t) = \mathcal{R}(X) + t$$

(R4) **Positive Homogeneity:** For all $X \in L^p(\Omega, \mathcal{F}, \mathbb{P})$ and $t \geq 0$,

$$\mathcal{R}(tX) = t\mathcal{R}(X)$$

Ph. Artzner, F. Delbaen, J.-M. Eber & D. Heath, *Coherent measures of risk.* Math. Finance, 1999.

Coherent Risk Measures

Some Good and *Not* So Good Properties?

Biconjugate Representation: (Recall $\mathcal{R}^*(\vartheta) = \sup_X\{\mathbb{E}[\vartheta X] - \mathcal{R}(X)\}$)

- $\mathcal{R}$ is proper, **convex** and lsc $\iff$

$$\mathcal{R}(X) = \sup\left\{\mathbb{E}[\vartheta X] - \mathcal{R}^*(\vartheta) \;\middle|\; \vartheta \in \mathrm{dom}(\mathcal{R}^*)\right\}.$$

- $\mathcal{R}$ is **translation equivariant** and **monotonic** $\iff$

$$\mathrm{dom}(\mathcal{R}^*) \subseteq \{\vartheta \in (L^p(\Omega, \mathcal{F}, \mathbb{P}))^* \mid \mathbb{E}[\vartheta] = 1,\ \vartheta \geq 0 \text{ a.s.}\}$$

- $\mathcal{R}$ is **positive homogeneous** $\iff$

$$\mathcal{R}(X) = \sup_{\vartheta \in \mathrm{dom}(\mathcal{R}^*)} \mathbb{E}[\vartheta X].$$

Optimal $\vartheta^\star \in \mathrm{dom}(\mathcal{R}^*)$ are called *risk identifiers*

Example (Conditional Value-at-Risk (CVaR)): $\mathcal{R}(X) = \frac{1}{1-\beta}\int_\beta^1 q_X(\beta)\,\mathrm{d}\beta$

$$\mathrm{dom}(\mathcal{R}^*) = \left\{\vartheta \in (L^p(\Omega, \mathcal{F}, \mathbb{P}))^* \,|\, \mathbb{E}[\vartheta] = 1,\ 0 \leq \vartheta \leq \frac{1}{1-\beta} \text{ a.s.}\right\}.$$

Differentiability: If $\mathcal{R} : L^p(\Omega, \mathcal{F}, \mathbb{P}) \to \mathbb{R}$ is **coherent**, then $\mathcal{R}$ is **Fréchet differentiable** $\iff$ $\exists\, \vartheta \in (L^p(\Omega, \mathcal{F}, \mathbb{P}))^*$ with $\vartheta \geq 0$ a.s., $\mathbb{E}[\vartheta] = 1$, and $\mathcal{R}(X) = \mathbb{E}[\vartheta X]$ for all $X \in L^p(\Omega, \mathcal{F}, \mathbb{P})$.

CVaR and Kusuoka Representation

Let $F_X(x) = \mathbb{P}(X \leq x)$, then CVaR is

$$\mathrm{CVaR}_\beta(X) := \frac{1}{1-\beta} \int_\beta^1 F_X^{-1}(\alpha)\, \mathrm{d}\alpha$$



In fact, all **law-invariant coherent** risk measures have the representation

$$\mathcal{R}(X) = \sup_{\mu \in \mathfrak{M}} \int_0^1 \mathrm{CVaR}_\beta(X)\, \mathrm{d}\mu(\beta)$$

where $\mathfrak{M}$ is a set of **probability measures** on $[0, 1]$.

Spectral Risk Measures: Given a probability measure ν on $[0, 1]$,

$$\begin{aligned} \mathcal{R}(X) &= \int_0^1 \mathrm{CVaR}_\beta(X)\, \mathrm{d}\nu(\beta) \\ &= \int_0^1 h(\beta) F_X^{-1}(\beta)\, \mathrm{d}\beta \quad \text{where} \quad h(\beta) := \int_0^\beta \frac{1}{1-\alpha}\, \mathrm{d}\nu(\alpha) \end{aligned}$$

S. Kusuoka, *On law-invariant coherent risk measures*, Advances in Math. Econ., 2001.

Risk Measure Examples

Risk Neutral:

$$\mathcal{R}(X) = \mathbb{E}[X]$$

is **law invariant** and **coherent**.

Mean-Plus-Deviation:

$$\mathcal{R}(X) = \mathbb{E}[X] + c\mathbb{E}[|X - \mathbb{E}[X]|^p]^{1/p}, \quad c > 0$$

is **law invariant** and satisfies (R1), (R3) and (R4), but **not** (R2).

Mean-Plus-Upper-Semideviation:

$$\mathcal{R}(X) = \mathbb{E}[X] + c\mathbb{E}[(X - \mathbb{E}[X])_+^p]^{1/p}, \quad c \in [0, 1]$$

is **law invariant** and **coherent**.

Conditional Value-at-Risk:

$$\mathcal{R}(X) = \frac{1}{1-\beta}\int_\beta^1 F_X^{-1}(\eta)\,\mathrm{d}\eta = \inf_{t\in\mathbb{R}}\left\{t + \frac{1}{1-\beta}\mathbb{E}[(X-t)_+]\right\}, \quad 0 \le \beta < 1$$

is **law invariant** and **coherent**.

Entropic Risk:

$$\mathcal{R}(X) = \lambda^{-1}\ln\mathbb{E}[\exp(\lambda X)], \ \lambda > 0$$

is **law invariant** and satisfies (R1), (R2) and (R3), but **not** (R4).

More Measures of Risk

One can quantify risk using the optimized certainty equivalent risk measure

$$\mathcal{R}(X) = \inf_{t \in \mathbb{R}} \{ t + \mathbb{E}[v(X - t)] \}$$

where $v : \mathbb{R} \to \mathbb{R}$ is a convex *regret* function that satisfies

$$v(0) = 0, \quad v(x) > x \quad \forall\, x \neq 0$$

Relation to Utility: $u(x) = -v(-x)$ is a *utility* function

Properties: $\mathcal{R}$ is convex and translation equivariant

$\mathcal{R}$ is **positive homogeneous** $\iff$ v is piecewise linear with kink at 0

$\mathcal{R}$ is **monotonic** $\iff$ v is nondecreasing

Mean-Plus-Variance



CVaR



Entropic Risk



A. Ben Tal & M. Teboulle, *An old-new concept of convex risk measures: The optimized certainty equivalents*, Math. Finance, 2007.

The Risk Quadrangle

$$\mathcal{R}(X) = \mathbb{E}[X] + \mathcal{D}(X) = \min_t \{t + \mathcal{V}(X - t)\}$$

$$\mathcal{V}(X) = \mathbb{E}[X] + \mathcal{E}(X)$$

Optimization



Estimation

$$\mathcal{D}(X) = \mathcal{R}(X) - \mathbb{E}[X] = \min_t \mathcal{E}(X - t)$$

$$\mathcal{E}(X) = \mathcal{V}(X) - \mathbb{E}[X]$$

- $\mathcal{R}$ quantifies **hazard** — Used in optimization as objective function or constraint
- $\mathcal{E}$ quantifies **nonzeroness** — Used in regression analysis, e.g., polynomial chaos
- $\mathcal{V}$ quantifies **displeasure** for positive values — Used to define risk via *disutility*
- $\mathcal{D}$ quantifies **nonconstancy** — Used to define risk via *variability*

Quantile Quadrangle: $0 < \alpha < 1$

$$\mathcal{R}(X) = \mathrm{CVaR}_\alpha(X) \qquad \mathcal{D}(X) = \mathrm{CVaR}_\alpha(X - \mathbb{E}[X])$$

$$\mathcal{V}(X) = \tfrac{1}{1-\alpha}\mathbb{E}[X_+] \qquad \mathcal{E}(X) = \mathbb{E}[\tfrac{\alpha}{1-\alpha}X_+ + X_-]$$

$$\mathcal{S}(X) = q_\alpha(X)$$

Safety Margins Quadrangle: $c > 0$

$$\mathcal{R}(X) = \mathbb{E}[X] + c\sigma(X) \qquad \mathcal{D}(X) = c\sigma(X)$$

$$\mathcal{V}(X) = \mathbb{E}[X] + c\|X\|_2 \qquad \mathcal{E}(X) = c\|X\|_2$$

$$\mathcal{S}(X) = \mathbb{E}[X]$$

R. T. Rockafellar & S. Uryasev, *The fundamental risk quadrangle in risk management, optimization, and statistical estimation*, Surveys in OR & Managment Science, 2013.

Superquantile Quadrangle

Choosing the **uniform** probability measure on $[\beta, 1]$,

$$\nu(S) = \frac{1}{1-\beta} \int_S \mathbb{1}_{[\beta,1]}(\alpha)\, \mathrm{d}\alpha,$$

produces the **second-order** CVaR

$$\mathcal{R}(X) = \frac{1}{1-\beta} \int_\beta^1 \mathrm{CVaR}_\alpha(X)\, \mathrm{d}\alpha$$



Second-order CVaR is a product of the **risk quadrangle**:

$$\mathcal{R}(X) = \frac{1}{1-\beta} \int_\beta^1 \mathrm{CVaR}_\alpha(X)\, \mathrm{d}\alpha \qquad \mathcal{D}(X) = \frac{1}{1-\beta} \int_\beta^1 \mathrm{CVaR}_\alpha(X - \mathbb{E}[X])\, \mathrm{d}\alpha$$

$$\mathcal{V}(X) = \frac{1}{1-\beta} \int_0^1 (\mathrm{CVaR}_\alpha(X))_+\, \mathrm{d}\alpha \qquad \mathcal{E}(X) = \frac{1}{1-\beta} \int_0^1 (\mathrm{CVaR}_\alpha(X))_+\, \mathrm{d}\alpha - \mathbb{E}[X]$$

$$\mathcal{S}(X) = \mathrm{CVaR}_\beta(X)$$

R. T. Rockafellar & J. O. Royset, *Random variables, monotone relations, and convex analysis*, Math. Programming, 2014.

Example — CVaR

Optimal Control of 1D Elliptic Equation

Let $\gamma = 10$, $D = (-1, 1)$, and $w \equiv 1$ and consider

$$\underset{z \in L^2(-1,1)}{\text{minimize}} \quad J(z) = \frac{1}{2}\text{CVaR}_\beta \left[\int_{-1}^{1} (S(z)(\cdot, x) - 1)^2 \, \mathrm{d}x \right] + \frac{\gamma}{2} \int_{-1}^{1} z(x)^2 \, \mathrm{d}x$$

where $S(z) = u \in L^2(\Omega, \mathcal{F}, \mathbb{P}; H_0^1(-1, 1))$ solves the weak form of

$$\begin{aligned} -\partial_x \left(\epsilon(\omega, x) \partial_x u(\omega, x) \right) &= f(\omega, x) + z(x) && x \in D, \text{ a.s.}, \\ u(\omega, -1) = 0, \quad u(\omega, 1) &= 0 && \text{a.s.} \end{aligned}$$

$\Omega = [-0.1, 0.1] \times [-0.5, 0.5]$ is endowed with the uniform density, and the random field coefficients are

$$\epsilon(\omega, x) = 0.1 \cdot \mathbb{1}_{(-1, \omega_1)} + 10 \cdot \mathbb{1}_{(\omega_1, 1)}, \quad \text{and} \quad f(\omega, x) = \exp(-(x - \omega_2)^2).$$

Example — CVaR

Sample Approximation: Monte Carlo with 10,000 samples.



$\vartheta^* = 0$ and $\vartheta^* = (1-\beta)^{-1}$

Outline

Motivating Applications

General Problem Formulation

Quantifying Risk

Probabilistic Optimization

What if our uncertainty is uncertain?

Computational Solution Methods

Software

References

Probabilistic Hazard

Standard Engineering Prospective

$X = \widehat{J}(z) =$ "cost" signaling "danger"



Probability of failure: $\mathcal{R}(X) = p_x(X) = \mathbb{P}(X > x)$

- How to compute or at least estimate?
- How to cope with control variables z in optimization?
  Both $p_x(X)$ and the threshold change with z!

Troubles with this concept:

- Poor mathematical behavior is a serious handicap.
- Failure probability ignores the **degree** of failure.

Buffered Probabilities

Rockafellar & Royset (2013), Mafusalov & Uryasev (2014), Norton & Uryasev (2014)

Utilizing **CVaR** in place of **quantile** in reliability



Buffered probability of failure: $\mathcal{R}(X) = \bar{p}_x(X) = \mathbb{P}(X > \tau(x))$
where $\tau(x)$ is determined by $\mathrm{CVaR}_{(1-\bar{p}_x(X))}(X) = \mathbb{E}[\, X \,|\, X > \tau(x)\,] = x.$

$$\mathrm{bPOE}_x[X] = 1 - \alpha \quad \text{where } \alpha \text{ solves} \quad \mathrm{CVaR}_\alpha[X] = x.$$





Buffered Probability Properties

- Optimization representation:

$$\mathrm{bPOE}_x[X] = \min_{t \geq 0} \mathbb{E}[(t(X - x) + 1)_+]$$

- Takes into account **values of outcomes in the distribution tail**
- Closed, quasi-convex and monotonic in random variable X
- Lowest quasi-convex (in X) upper bound of POE
- Continuous with respect to threshold $x \in [\mathbb{E}[X], \operatorname{ess\,sup} X)$
- Easy to manage (optimize with convex and linear programming)
- $$\mathrm{CVaR}_\alpha[X] \leq x \quad \Longleftrightarrow \quad \mathrm{bPOE}_x[X] \leq 1 - \alpha$$

Objective function in optimization representation is **nonsmooth**!

Question: Is it possible to account for higher-order tail moments?

Higher-Moment Coherent Risk Measures

Higher-Moment Coherent Risk (HMCR) measures with $p \geq 1$ and $\beta \in [0, 1)$

$$\mathrm{HMCR}_{p,\beta}[X] = \inf_{t \in \mathbb{R}} \left\{ t + \frac{1}{1-\beta} \mathbb{E}[(X-t)_+^p]^{1/p} \right\}$$

1. As the name suggests, $\mathrm{HMCR}_{p,\beta}$ is coherent and law invariant
2. When $p = 1$, we have that $\mathrm{HMCR}_{1,\beta}[X] = \mathrm{CVaR}_\beta[X]$
3. $\mathrm{HMCR}_{p,\beta}$ is generated from the risk quadrangle with regret measure

$$\mathcal{V}(X) = \frac{1}{1-\beta} \mathbb{E}[(X)_+^p]^{1/p}$$

Properties of HMCR: Suppose X is not degenerate (constant)

1. $p \mapsto \mathrm{HMCR}_{p,\beta}[X]$ is nondecreasing
2. $\beta \mapsto \mathrm{HMCR}_{p,\beta}[X]$ is nondecreasing and continuous
3. In fact, $\beta \mapsto \mathrm{HMCR}_{p,\beta}[X]$ is strictly increasing on $[0, 1 - \pi_X)$ with

$$\pi_X = \mathrm{prob}(X = \operatorname{ess\,sup} X)$$

4. $\mathrm{HMCR}_{p,0}[X] = \mathbb{E}[X]$ and $\mathrm{HMCR}_{p,1}[X] = \operatorname{ess\,sup} X$

$\beta \mapsto \mathrm{HMCR}_{p,\beta}[X]$ **has a nondecreasing and continuous inverse!**

P. A. Krokhmal, *Higher moment coherent risk measures*, Quantitative Finance, 2007.

Higher-Moment bPOE Properties

Kouri (2018)

- Optimization representation:

$$\mathrm{bPOE}_{p,x}[X] = \min_{t \geq 0} \mathbb{E}[(t(X-x)+1)_+^p]^{1/p}$$

- Takes into account **moments** of outcomes in the distribution tail
- Closed, quasi-convex and monotonic in random variable X
- Continuous with respect to threshold $x \in [\mathbb{E}[X], \operatorname{ess\,sup} X)$
- Objective function in optimization representation is **smooth** in X
- $$\mathrm{HMCR}_{p,\alpha}[X] \leq x \quad \Longleftrightarrow \quad \mathrm{bPOE}_{p,x}[X] \leq 1-\alpha$$
- $$\mathrm{bPOE}_x[X] \leq (\mathrm{bPOE}_{2,x}[X])^2 \leq \ldots \leq (\mathrm{bPOE}_{p,x}[X])^p$$

Example: Second-Moment Buffered Probability

Suppose $X \sim N(0,1)$ with cdf Φ and pdf ϕ. Let $x \geq 0$ then

$$Z := (t(X - x) + 1) \sim N(1 - tx, t) \quad \forall\, t > 0$$

Therefore, the buffered probability of X exceeding x is

$$\mathrm{bPOE}_x[X] = \min_{t \geq 0} \{(1 - tx)(1 - \Phi(x - 1/t)) + t\phi(x - 1/t)\}$$

and the second order buffered probability of X exceeding x is

$$(\mathrm{bPOE}_{2,x}[X])^2 = \min_{t \geq 0} \{((1-tx)^2 + t^2)(1 - \Phi(x - 1/t)) + t(1 - tx)\phi(x - 1/t)\}.$$

x	$\mathrm{POE}_x[X]$	$\mathrm{bPOE}_x[X]$	$(\mathrm{bPOE}_{2,x}[X])^2$	$\mathrm{bPOE}_{2,x}[X]$
0	0.5	1	1	1
1	0.15866	0.38109	0.46235	0.67996
2	0.02275	0.05799	0.07423	0.27246
3	0.00135	0.00353	0.00463	0.06806
4	0.00003	0.00008	0.00011	0.01056

$x = 2$



Order 1 and Order 2

3D Topology Optimization with Buffered Probability

Given compliance tolerance c_0, probability $p_0 \in (0,1)$, order $q \geq 1$,

$$\min_{0 \leq z \leq 1} \int_D z \, \mathrm{d}x =: \mathrm{vol}(z) \quad \text{subject to} \quad \mathrm{bPOE}_{q,c_0}\left(\int_D F \cdot S(z) \mathrm{d}x\right) \leq 1 - p_0$$

where $S(z) = u$ solves the **linear elasticity equations**

$$\begin{aligned} -\nabla \cdot (\mathbf{E}(z) : \varepsilon u) &= F, && \text{in } D \\ \varepsilon u &= \tfrac{1}{2}(\nabla u + \nabla u^\top), && \text{in } D \\ u &= 0, && \text{on } \Gamma_D \\ \varepsilon u : n &= 0, && \text{on } \partial D \setminus \Gamma_D \end{aligned}$$



Numerical Results

Spatial Discretization: Q1 FEM on a uniform $32 \times 16 \times 16$ mesh

Stochastic Discretization: $Q = 120$ Monte Carlo samples

Problem Data: $p_0 = 0.75$ and $c_0 = 2\mathbb{E}\left[\int_D F \cdot S(\mathbf{1})\mathrm{d}x\right]$

Mean Value



Risk Neutral



bPOE



	MV	RN	bPOE
Volume Fraction	49.061%	47.634%	67.204%

Numerical Results

Spatial Discretization: Q1 FEM on a uniform $32 \times 16 \times 16$ mesh

Stochastic Discretization: $Q = 120$ Monte Carlo samples

Problem Data: $p_0 = 0.75$ and $c_0 = 2\mathbb{E}\left[\int_D F \cdot S(\mathbf{1})\mathrm{d}x\right]$

Mean Value **Risk Neutral** **bPOE**



Topology changes from beam to shell!

	MV	RN	bPOE
Volume Fraction	49.061%	47.634%	67.204%

Numerical Results

Spatial Discretization: Q1 FEM on a uniform $32 \times 16 \times 16$ mesh

Stochastic Discretization: $Q = 120$ Monte Carlo samples

Problem Data: $p_0 = 0.75$ and $c_0 = 2\mathbb{E}\left[\int_D F \cdot S(\mathbf{1})\mathrm{d}x\right]$

bPOE_{c_0}



bPOE_{2,c_0}



bPOE_{3,c_0}



Order	1	2	3
Volume Fraction	67.204%	77.369%	80.075%

Numerical Results

Spatial Discretization: Q1 FEM on a uniform $32 \times 16 \times 16$ mesh

Stochastic Discretization: $Q = 120$ Monte Carlo samples

Problem Data: $p_0 = 0.75$ and $c_0 = 2\mathbb{E}\left[\int_D F \cdot S(\mathbf{1})\mathrm{d}x\right]$

bPOE_{c_0} bPOE_{2,c_0} bPOE_{3,c_0}



Topology changes from beam to shell!

Order	1	2	3
Volume Fraction	67.204%	77.369%	80.075%

Outline

Motivating Applications

General Problem Formulation

Quantifying Risk

Probabilistic Optimization

What if our uncertainty is uncertain?

Computational Solution Methods

Software

References

What if our uncertainty is uncertain?

Distributionally Robust Stochastic Programming

$(\Omega, \mathcal{F})$ is a measurable space and prob. measure is *unknown*.
Consider

$$\min_{z \in Z_{\text{ad}}} \mathcal{R}(\widehat{J}(z)) = \sup_{P \in \mathfrak{A}} \mathbb{E}_P[\widehat{J}(z)].$$

Ambiguity Set: $\mathfrak{A} \subset \{P : \mathcal{F} \to [0,1] \mid P(\Omega) = 1\}$ defined by data.
For example:

- **Moment Matching:** Given generalized moment data $m_1, \ldots, m_N$,

$$\mathfrak{A} = \{P : \mathcal{F} \to [0,1] \mid P(\Omega) = 1,\ \mathbb{E}_P[\psi_i] = m_i,\ i = 1, \ldots, N\}.$$

- **Φ-Divergence (e.g., Kullback-Leibler, χ^2, TV, Hellinger, ...):** Given a nominal P_0 and $\epsilon > 0$,

$$\mathfrak{A} = \{P : \mathcal{F} \to [0,1] \mid P(\Omega) = 1,\ D_\Phi(P, P_0) \leq \epsilon\}.$$

- **Wasserstein Distance:** Given a nominal P_0 and $\epsilon > 0$,

$$\mathfrak{A} = \left\{ P : \mathcal{F} \to [0,1] \mid P(\Omega) = 1,\ \sup_{f \in \mathcal{L}} \int_\Omega f(\omega)\, \mathrm{d}(P - P_0)(\omega) \leq \epsilon \right\}.$$

Example: Moment Matching

Let $\psi_i : \Omega \to \mathbb{R}$ be $\mathcal{F}$-measurable functions and $m_i \in \mathbb{R}$ for $i = 1, \ldots, N$

$$\mathfrak{A} = \left\{ P : \mathcal{F} \to [0,1] \;\mid\; P(\Omega) = 1, \quad \begin{array}{l} \mathbb{E}_P[\psi_i] = m_i,\ i = 1, \ldots, N_e \\ \\ \mathbb{E}_P[\psi_i] \leq m_i,\ i = N_e + 1, \ldots, N \end{array} \right\}.$$

Theorem (Rogosinski): If $\mathfrak{A} \neq \emptyset$, then for each $z \in Z$ there exists ω_i and $p_i \geq 0$ with $p_1 + \cdots + p_{N+1} = 1$ such that

$$\mathcal{R}(\widehat{J}(z)) = \sup_{P \in \mathfrak{A}} \mathbb{E}_P[\widehat{J}(z)] = \sum_{i=1}^{N+1} p_i J([S(z)](\omega_i), z, \omega_i)$$

W. W. Rogosinski, *Moments of non-negative mass*, Proceedings of the Royal Society of London: Series A, Math. and Phys. Sciences, 1958.

Example: Φ-Divergence

Supppose

(i) A **nominal** probability measure P_0 is given,

(ii) The random variable $X \in L^p(\Omega, \mathcal{F}, P_0)$, and

(iii) $\Phi : \mathbb{R} \to [0, \infty]$ is convex lower semicontinuous satisfying

$$\Phi(1) = 0 \quad \text{and} \quad \Phi(x) = \infty \quad \forall\, x < 0.$$

Define, for fixed $\epsilon > 0$,

$$\mathfrak{A} = \{\vartheta \in (L^p(\Omega, \mathcal{F}, P_0))^* \mid \mathbb{E}_{P_0}[\vartheta] = 1,\ \vartheta \geq 0,\ \mathbb{E}_{P_0}[\Phi(\vartheta)] \leq \epsilon\}.$$

Then $\mathcal{R}(X) = \sup_{\vartheta \in \mathfrak{A}} \mathbb{E}_{P_0}[\vartheta X] = \inf_{\lambda \geq 0,\, \mu} \{\lambda\epsilon + \mu + \mathbb{E}_{P_0}[(\lambda\Phi)^*(X - \mu)]\}$

is a **law-invariant coherent** risk measure!

Example (Kullback-Leibler Divergence): $\Phi(x) = x\ln(x) - x + 1,\ x \geq 0$

$$\mathcal{R}(X) = \inf_{\lambda > 0} \left\{\lambda c + \lambda \ln \mathbb{E}_{P_0}\left[e^{X/\lambda}\right]\right\}.$$

A. Ben Tal & M. Teboulle, *Penalty functions and duality in stochastic programming via phi-divergence functionals*, Mathematics of Operations Research, 1987.

Robust Probabilitistic Optimization

Shapiro, Mafusalov, Uryasev, Kouri (2018)

When $\mathbb{P}$ is unknown, we can similarly *robustify* the POE and bPOE.

Probability: In general, we have that

$$\mathrm{POE}_x^\star(X) = \sup_{P\in\mathfrak{A}} P(X > x) = \sup_{P\in\mathfrak{A}} \mathbb{E}_P[\mathbb{1}_A] = \mathcal{R}(\mathbb{1}_A)$$

where $A = \{\omega \in \Omega \,|\, X(\omega) > x\}$ and $\mathcal{R}$ is a **coherent** risk measure!

Buffered Probability: Under mild regularity conditions, we have

$$\begin{aligned}\mathrm{bPOE}_x^\star(X) = \sup_{P\in\mathfrak{A}} \min_{t\geq 0} \mathbb{E}_P[(t(X-x)+1)_+] &= \min_{t\geq 0} \sup_{P\in\mathfrak{A}} \mathbb{E}_P[(t(X-x)+1)_+] \\ &= \min_{t\geq 0} \mathcal{R}((t(X-x)+1)_+)\end{aligned}$$

where $\mathcal{R}$ is a **coherent** risk measure! For Φ-divergence ambiguity,

$$\mathrm{bPOE}_x^\star(X) = \min_{t\geq 0,\lambda\geq 0,\mu} \left\{\lambda\epsilon + \mu + \mathbb{E}_{P_0}[(\lambda\Phi)^*((t(X-x)+1)_+ - \mu)]\right\}.$$

Distributionally Robust Contaminant Mitigation

Problem Description

Model contaminant spread by advection-diffusion on $D = (0,1)^2$.
Determine controls that mitigate the contaminant

$$\min_z \ \mathcal{R}\left(\frac{\kappa_s}{2}\int_D S(z)^2\,\mathrm{d}x\right) + \wp(z) \quad \text{subject to} \quad 0 \le z \le 1$$

where $S(z) = u : \Omega \to H^1(D)$ solves

$$\begin{aligned} -\nabla\cdot(\epsilon(\omega)\nabla u(\omega)) + V(\omega)\cdot\nabla u(\omega) &= f(\omega) - Bz, && \text{in } D, \text{ a.s.} \\ u(\omega) &= 0, && \text{on } \Gamma_d, \text{ a.s.} \\ \epsilon(\omega)\nabla u(\omega)\cdot n &= 0, && \text{on } \partial D \setminus \Gamma_d, \text{a.s.,} \end{aligned}$$

$$Bz = \sum_{k=1}^{9} z_k \exp\left(\frac{-\|x - p_k\|_2^2}{2\sigma^2}\right) \quad \text{and} \quad \wp(z) = \kappa_c\|z\|_1 = \kappa_c\sum_{k=1}^{9} z_k.$$

Control	1	2	3	4	5	6	7	8	9
x_1	0.25	0.50	0.75	0.25	0.50	0.75	0.25	0.50	0.75
x_2	0.25	0.25	0.25	0.50	0.50	0.50	0.75	0.75	0.75

Total of 37 random variables.

Risk-Averse Contaminant Mitigation

Nominal Distribution: $\xi_k(\omega) \sim U(-1,1)$ with $k = 1, \ldots, 37$

Diffusivity:

$$\log(c\,\epsilon(\omega,x) - 0.5) = 1 + \xi_1(\omega)\left(\frac{\sqrt{\pi}L_c}{2}\right)^{1/2} + \sum_{n=2}^{10} \zeta_k \phi_k(x)\xi_k(\omega)$$

Advection:





Source:

$$f(\omega,x) = \sum_{k=1}^{5} \xi_{8+5k}(\omega) \exp\left(\frac{-(x_1 - \xi_{9+5k}(\omega))^2}{2\xi_{10+5k}(\omega)^2}\right) \exp\left(\frac{-(x_2 - \xi_{11+5k}(\omega))^2}{2\xi_{12+5k}(\omega)^2}\right).$$

Numerical Results

DRO with KL-Divergence Ambiguity

$$\mathcal{R}(X) = \inf_{\lambda>0} \left\{ \lambda c + \lambda \ln \mathbb{E}\left[e^{X/\lambda} \right] \right\}$$





$\log_{10}(c)$	1	2	3	4	5	6	7	8	9	obj
10^{-3}	—	0.410	—	—	1.000	—	—	—	—	3.465
10^{-2}	—	0.560	—	—	1.000	—	—	—	—	3.637
10^{-1}	—	1.000	—	—	1.000	—	—	—	—	4.186
1	—	1.000	—	0.580	1.000	0.709	—	—	—	5.939
5	1.000	1.000	0.249	1.000	1.000	1.000	—	—	—	8.124

Outline

Motivating Applications

General Problem Formulation

Quantifying Risk

Probabilistic Optimization

What if our uncertainty is uncertain?

Computational Solution Methods

Software

References

Methods for Stochastic Optimization

Minimizing the Expectation

1. **Stochastic Approximation (SA):** Stochastic subgradient descent only requires a single sample at every iteration.
2. **Progressive Hedging:** Decoupled deterministic optimization via alternating directions method of multipliers (ADMM).
3. **Sample Average Approximation (SAA):** (Quasi) Monte Carlo approximation of expected value.
4. **Adaptive Stochastic Collocation:** Deterministic quadrature approximation of expected value. Adaptivity using trust regions.

Note: The convergence of SA and SAA is **probabilistic**!

Note: Risk measures and probabilistic functionals are often nonsmooth $\implies$ polynomial approximation and derivative-based optimization may not apply.

The Finite Noise Assumption

Suppose there exists a random vector $\xi : \Omega \to \Xi \subseteq \mathbb{R}^M$ and functions $\bar{J} : U \times Z \times \Xi \to \mathbb{R}$ and $\bar{c} : U \times Z \times \Xi \to Y$ such that

$$J(u, z, \omega) = \bar{J}(u, z, \xi(\omega)) \quad \text{and} \quad c(u, z, \omega) = \bar{c}(u, z, \xi(\omega)).$$

Moreover, assume the probability law $\mathbb{P} \circ \xi^{-1}$ has Lebesgue density $\rho : \Xi \to \mathbb{R}$, i.e., $\mathrm{d}\mathbb{P} \circ \xi^{-1} = \rho \, \mathrm{d}\xi$.

This permits the change of variables from $\omega \in \Omega$ to $\xi \in \Xi$.
Analysis now performed in weighted Lebesgue space

$$L^p_\rho(\Xi) = \left\{ v : \Xi \to \mathbb{R} \mid \int_\Xi |v(\xi)|^p \rho(\xi) \, \mathrm{d}\xi < \infty \right\}.$$

$L^\infty_\rho(\Xi)$ and $L^p_\rho(\Xi; W)$ are similarly defined.

Independence: For adaptive stochastic collocation, we will assume that the components of ξ are independent and

$$\Xi = [a_1, b_1] \times \cdots \times [a_M, b_M] \quad \text{and} \quad \rho = \rho_1 \otimes \cdots \otimes \rho_M.$$

Stochastic Approximation

Set $\widehat{J}(z) = \bar{J}(u(z), z, \cdot)$. Let Z be Hilbert and Z_{ad} be closed convex. Given $z_k \in Z_{\text{ad}}$ and $G(z_k, \xi) = G_k(\xi)$ such that $\mathbb{E}[G_k(\xi)] \in \partial\mathbb{E}[\widehat{J}(z_k)]$, the SA iteration is

$$z_{k+1} = \Pi_{Z_{\text{ad}}}(z_k - \mu_k G_k(\xi_k)), \quad \mu_k > 0,$$

where ξ_k for $k = 1, \ldots$ is an iid sequence of realizations and

$$\Pi_{Z_{\text{ad}}}(z) = \arg\min_{\zeta \in Z_{\text{ad}}} \|z - \zeta\|_Z.$$

Note: $\Pi_{Z_{\text{ad}}}$ is (firmly) nonexpansive.

Note: For PDE-constrained optimization, SA requires a single **deterministic** state and adjoint solve per iteration!

Must solve:

$$\bar{c}(u, z_k, \xi_k) = 0 \quad \text{and} \quad \bar{c}_u(u_k, z_k, \xi_k)^* \lambda = -\bar{J}_u(u_k, z_k, \xi_k).$$

H. Robbins & S. Monro, *A stochastic approximation method*, An. Math. Statist., 1951.

Analysis for Linear-Elliptic Quadratic Control

Recall: **(Spaces)** $Z = L^2(D)$ and $Z_{\text{ad}} = \{z \in Z \mid z_a \leq z \leq z_b\}$, $U = H_0^1(D)$, $Y = H^{-1}(D)$, W is a Hilbert space such that $U \hookrightarrow W$,

$$\bar{J}(u, z, \xi) = \frac{1}{2}\|\mathbf{C}u - w\|_W^2 + \frac{\gamma}{2}\|z\|_Z^2$$

where $\mathbf{C} \in \mathcal{L}(U, W)$, $w \in W$ and $\gamma > 0$, and for $v \in U$

$$\langle \bar{c}(u, z, \xi), v\rangle_{-1,1} = \int_D (\overline{A}(\xi)\nabla u(x)) \cdot \nabla v(x)\,\mathrm{d}x - \int_D z(x)v(x)\,\mathrm{d}x.$$

Note: $\mathcal{J}(z) = \mathbb{E}[\widehat{J}(z)]$ is strongly convex with constant γ.

Stochastic Approximation: Given $z_k \in Z$ and $u_k \in U$ that solves $\bar{c}(u_k, z_k, \xi_k) = 0$

$$G_k(\xi_k) = \gamma z_k + \lambda_k$$

where λ_k solves the adjoint equation

$$\int_D (\overline{A}(\xi_k)\nabla \lambda_k(x)) \cdot \nabla v(x)\,\mathrm{d}x = -\langle \mathbf{C}u_k - w, \mathbf{C}v\rangle_W \quad \forall\, v \in U.$$

Analysis for Linear-Elliptic Quadratic Control

Let $z^* \in Z_{\text{ad}}$ minimize $\mathcal{J}(z)$ over Z_{ad} then (since $\Pi_{Z_{\text{ad}}}$ is nonexpansive)

$$\begin{aligned}\mathbb{E}[\|z_{k+1} - z^*\|_Z^2] &= \mathbb{E}[\|\Pi_{Z_{\text{ad}}}(z_k - \mu_k G_k(\xi_k)) - \Pi_{Z_{\text{ad}}}(z^*)\|_Z^2] \\ &\leq \mathbb{E}[\|z_k - z^*\|_Z^2] + \mu_k^2 \mathbb{E}[\|G_k(\xi_k)\|_Z^2] - 2\mu_k \mathbb{E}[\langle G_k(\xi_k), z_k - z^*\rangle_Z]\end{aligned}$$

z_k only depends on $\xi_1, \ldots, \xi_{k-1}$ (which are iid), thus

$$\begin{aligned}\mathbb{E}[\langle z_k - z^*, G_k(\xi_k)\rangle_Z] &= \mathbb{E}[\mathbb{E}[\langle z_k - z^*, G_k(\xi_k)\rangle_Z | \xi_1, \ldots, \xi_{k-1}]] && \text{Law of Total Exp.} \\ &= \mathbb{E}[\langle z_k - z^*, \mathbb{E}[G_k(\xi_k)|\xi_1, \ldots, \xi_{k-1}]\rangle_Z] && \text{Fubini's Theorem} \\ &= \mathbb{E}[\langle z_k - z^*, \mathbb{E}[\nabla \overline{F}(z_k)]\rangle_Z] \\ &\geq \mathbb{E}[\langle z_k - z^*, \mathbb{E}[\nabla \overline{F}(z_k) - \nabla \overline{F}(z^*)]\rangle_Z && \text{Optimality of } z^* \\ &\geq \gamma \mathbb{E}[\|z_k - z^*\|_Z^2]. && \text{Strong Convexity of } \mathcal{J}\end{aligned}$$

Analysis for Linear-Elliptic Quadratic Control

Since Z_{ad} is bounded and u, λ depend continuously on z, we have

$$\mathbb{E}[\|G(z,\xi)\|_Z^2] \leq M^2 \quad \forall z \in Z_{\text{ad}}$$

$$\implies \quad \mathbb{E}[\|z_{k+1} - z^*\|_Z^2] \leq \mathbb{E}[\|z_k - z^*\|_Z^2] + \mu_k^2 M^2 - 2\mu_k \gamma \mathbb{E}[\|z_k - z^*\|_Z^2].$$

Now, set $\mu_k = \theta/k$, then

$$\mathbb{E}[\|z_{k+1} - z^*\|_Z^2] \leq \left(1 - \frac{2\gamma\theta}{k}\right) \mathbb{E}[\|z_k - z^*\|_Z^2] + \frac{\theta^2 M^2}{k^2} \qquad \text{Previous Results}$$

$$\leq \frac{\max\{\theta^2 M^2 (2\gamma\theta - 1)^{-1}, \|z_1 - z^*\|_Z^2\}}{k}. \qquad \text{Use Induction}$$

Minimizing the right hand side with $\theta > 0$ gives $\theta^* = 1/\gamma$.

Note: The expected decay at each iteration is $\mathcal{O}(k^{-1})$

$\implies$ to reach tolerance ε requires $\mathcal{O}(\varepsilon^{-1})$ iterations (on average)!

Progressive Hedging

Problem Assumptions: Suppose $\widehat{J}(\cdot, \xi)$ is a **convex random loss** and ξ is discretely distributed. Consider the convex program

$$\underset{z \in Z_{\text{ad}}}{\text{Minimize}} \left\{ \mathbb{E}[\widehat{J}(z, \xi)] = \sum_{k=1}^{N} p_k \widehat{J}(z, \xi_k) \right\}.$$

Progressive Hedging Algorithm:

Given $\hat{z} \in Z$ and a Z-valued r.v. $W(\xi)$ with $\mathbb{E}[W(\xi)] = 0$.

1. Compute $\zeta(\xi) \in Z_{\text{ad}}$ a.s. that approximately solves

$$\underset{z \in Z_{\text{ad}}}{\text{Minimize}} \left\{ \widehat{J}(z, \xi) + \langle W(\xi), z \rangle_Z + \tfrac{r}{2} \|z - \hat{z}\|_Z^2 \right\} \quad \text{a.s.}$$

2. Update $\hat{z} = \mathbb{E}[\zeta(\xi)] = p_1 \zeta_1 + \ldots + p_N \zeta_N$.
3. Update $W(\xi) = W(\xi) + r(\zeta(\xi) - \hat{z})$.

Step 1 requires solving decoupled deterministic convex opt. problems!

However, objective function $\widehat{J}(\cdot, \xi)$ must be convex . . .

R. T. Rockafellar & R. J.-B. Wets, *Scenarios and policy aggregation in optimization under uncertainty*, Math. Oper. Res., 1991.

Sample Average Approximation

Idea: Approximate expected value in $\mathcal{J}$ using Monte Carlo.

Let $\xi_1, \ldots, \xi_N$ be iid random samples of ξ, then solve

$$\underset{z \in Z_{\text{ad}}}{\text{Minimize}} \left\{ \widehat{\mathcal{J}}_N(z) = \frac{1}{N} \sum_{k=1}^{N} \widehat{J}(z, \xi_k) \right\}.$$

Apply nonlinear programming algorithms to solve numerically.

Linear-Elliptic Quadratic Control:

(i) Let $z_N^* \in Z_{\text{ad}}$ minimize $\widehat{\mathcal{J}}_N$ over Z_{ad}

(ii) Let $z^* \in Z_{\text{ad}}$ minimize $\mathcal{J}$ over Z_{ad}.

Strong convexity of $\mathcal{J}$ and optimality of z_N^*, z^* imply

$$\begin{aligned} \gamma \|z_N^* - z^*\|_Z^2 &\leq \langle z_N^* - z^*, \nabla \mathcal{J}(z_N^*) - \nabla \mathcal{J}(z^*) \rangle_Z \\ &\leq \langle z_N^* - z^*, \nabla \mathcal{J}(z_N^*) - \nabla \widehat{\mathcal{J}}_N(z_N^*) \rangle_Z \end{aligned}$$

Therefore, $\gamma \|z^* - z_N^*\|_Z \leq \left\| \mathbb{E}[\lambda] - \frac{1}{N} \sum_{k=1}^{N} \lambda_k \right\|_Z$ "$=$" $\mathcal{O}(N^{-\frac{1}{2}})$ **Probabilistic!**

Stochastic Collocation

Idea: Approximate expected value in $\mathcal{J}$ using quadrature.

Let $\xi_1, \ldots, \xi_N$ be quad. points with weights $w_1, \ldots, w_N$, then solve

$$\underset{z \in Z_{\text{ad}}}{\text{Minimize}} \left\{ \widehat{\mathcal{J}}_N(z) = \sum_{k=1}^{N} w_k \widehat{J}(z, \xi_k) \right\}.$$

Apply nonlinear programming algorithms to solve numerically.

Linear-Elliptic Quadratic Control:

(i) Let $z_N^* \in Z_{\text{ad}}$ minimize $\widehat{\mathcal{J}}_N$ over Z_{ad}

(ii) Let $z^* \in Z_{\text{ad}}$ minimize $\mathcal{J}$ over Z_{ad}.

Strong convexity of $\mathcal{J}$ and optimality of z_N^*, z^* imply

$$\begin{aligned} \gamma \|z_N^* - z^*\|_Z^2 &\leq \langle z_N^* - z^*, \nabla \mathcal{J}(z_N^*) - \nabla \mathcal{J}(z^*) \rangle_Z \\ &\leq \langle z_N^* - z^*, \nabla \mathcal{J}(z_N^*) - \nabla \widehat{\mathcal{J}}_N(z_N^*) \rangle_Z \end{aligned}$$

Therefore, $\gamma \|z^* - z_N^*\|_Z \leq \left\| \mathbb{E}[\lambda] - \sum_{k=1}^{N} w_k \lambda_k \right\|_Z =$ **Quad. Error**

Sparse Grids and Adaptivity

Gerstner and Griebel 2003

- **1D Operators:** For $k = 1, \ldots, M$, $\mathbb{E}_k^0 \equiv 0$ and

$$\Delta_k^i \equiv \mathbb{E}_k^i - \mathbb{E}_k^{i-1} \quad \text{where} \quad \mathbb{E}_k^i(g) \xrightarrow{i \to \infty} \int_{\Xi_k} \rho_k(\xi) g(\xi) \mathrm{d}\xi$$

- **Sparse-Grid Operator:** For an index set $\mathcal{I} \subset \mathbb{N}^M$,

$$\mathbb{E}_{\mathcal{I}} \equiv \sum_{\mathbf{i} \in \mathcal{I}} (\Delta_1^{i_1} \otimes \cdots \otimes \Delta_M^{i_M})$$

- **Admissibility:** $\mathbf{i} \in \mathcal{I}$ and $\mathbf{i} \geq \mathbf{j} \implies \mathbf{j} \in \mathcal{I}$
- **Error:** Given the index set $\mathcal{I} \subset \mathbb{N}^M$, the error is

$$\mathbb{E} - \mathbb{E}_{\mathcal{I}} = \sum_{\mathbf{i} \notin \mathcal{I}} (\Delta_1^{i_1} \otimes \cdots \otimes \Delta_M^{i_M})$$

- **Adaptivity:** Pick $\mathbf{i} \notin \mathcal{I}$ s.t. $\mathcal{I} \cup \{\mathbf{i}\}$ admissible and $\Delta_1^{i_1} \otimes \cdots \otimes \Delta_M^{i_M}$ "large"



Trust-Region Algorithm

Given: z_0, $m_0(s) \approx \mathcal{J}(z_0 + s)$, $\mathcal{J}_0 \approx \mathcal{J}$, $\Delta_0 \geq 0$, and $\mathrm{gtol} > 0$.
While $\|\nabla m_k(s)\|_{\mathcal{Z}} > \mathrm{gtol}$

1. **Model Update:** Choose a new $m_k(s) \approx \mathcal{J}(z_k + s)$. $\leftarrow$ **ADAPTIVITY**
2. **Step Computation:** Approximate a solution, s_k, to the subproblem

$$\min_{s \in \mathcal{Z}} \; m_k(s) \quad \text{subject to} \quad \|s\|_{\mathcal{Z}} \leq \Delta_k.$$

3. **Objective Update:** Choose a new $\mathcal{J}_k(z) \approx \mathcal{J}(z)$. $\leftarrow$ **ADAPTIVITY**
4. **Step Acceptance:** Compute

$$\rho_k = \frac{\mathcal{J}_k(z_k) - \mathcal{J}_k(z_k + s_k)}{m_k(0) - m_k(s_k)}.$$

   If $\rho_k \geq \eta \in (0, 1)$, then $z_{k+1} = z_k + s_k$ else $z_{k+1} = z_k$.
5. **Trust Region Update:** Choose a new trust region radius, Δ_{k+1}.

EndWhile

Inexact Gradients and Objective Functions

Kouri, Heinkenschloss, Ridzal, and van Bloemen Waanders (2013, 2014)

Inexact Gradients

There exists $c > 0$ independent of k such that

$$\|\nabla m_k(0) - \nabla \mathcal{J}(z_k)\|_{\mathcal{Z}} \leq c \min\{\|\nabla m_k(0)\|_{\mathcal{Z}}, \Delta_k\}$$

(Carter 1989, Heinkenschloss and Vicente 2001).

Inexact Objective Functions

There exists $K > 0$, $\omega \in (0,1)$, and $\theta(z,s) \to 0$ as $r \to 0$ such that

$$|(\mathcal{J}(z_k) - \mathcal{J}(z_k + s_k)) - (\mathcal{J}_k(z_k) - \mathcal{J}_k(z_k + s_k))| \leq K\theta(z_k, s_k)$$
$$\theta(z_k, s_k)^{\omega} \leq \eta \min\left\{(m_k(0) - m_k(s_k)), r_k\right\}.$$

Here, $\eta > 0$ is tied to algorithmic parameters and $\lim_{k\to\infty} r_k = 0$.
(Carter 1989, Ziems and Ulbrich 2013).

- **Cannot** compute $\mathcal{J}(z_k)$ and $\nabla \mathcal{J}(z_k)$;
- Control *a posteriori* errors using **adaptive sparse grids**.

Optimal Control of Steady Burger's Equation

Let $\gamma = 10^{-3}$, $\Omega_o = \Omega_c = \Omega = (0,1)$, and $w \equiv 1$ and consider

$$\min_{z \in L^2(0,1)} \quad \mathcal{J}(z) = \frac{1}{2}\mathbb{E}\left[\int_0^1 (u(\cdot,x;z)-1)^2 \, \mathrm{d}x\right] + \frac{\gamma}{2}\int_0^1 z(x)^2 \, \mathrm{d}x$$

where $u = S(z) \in L^3_\rho(\Xi; H^1(0,1))$ solves the weak form of

$$\begin{aligned} -\nu(\xi)\partial_{xx}u(\xi,x) + u(\xi,x)\partial_x u(\xi,x) &= f(\xi,x) + z(x) \qquad && (\xi,x) \in \Xi \times \Omega, \\ u(\xi,0) = d_0(\xi), \quad u(\xi,1) &= d_1(\xi) && \xi \in \Xi. \end{aligned}$$

$\Xi = [-1,1]^4$ is endowed with the uniform density $\rho(\xi) \equiv 2^{-4}$, and the random field coefficients are

$$\nu(\xi) = 10^{\xi_1 - 2}, \quad f(\xi,x) = \frac{\xi_2}{100}, \quad d_0(\xi) = 1 + \frac{\xi_3}{1000}, \quad \text{and} \quad d_1(\xi) = \frac{\xi_4}{1000}.$$

Adaptive Sparse Grid Results

Spatial: Piecewise Linear Finite Elements
Stochastic: Maximum Level 8 Clenshaw-Curtis Sparse Grids

Algorithm	NonlinPDE	CP_{obj}	LinearPDE	CP_{grad}	Rel. Err.
Newton-CG	45,224 (1.0)	7,537	489,906 (1.0)	7,537	–
Grad. Adapt.	45,531 (1.0)	7,537	3,405 (143.9)	249	2.89×10^{-6}
Full Adapt.	603 (75.0)	23	3,405 (143.9)	249	2.89×10^{-6}



Outline

Motivating Applications

General Problem Formulation

Quantifying Risk

Probabilistic Optimization

What if our uncertainty is uncertain?

Computational Solution Methods

Software

References

The Rapid Optimization Library

RAPID OPTIMIZATION LIBRARY

Stochastic Optimization Capabilities

1. **Risk Measures:** CVaR, Expectation, Mean-Plus-(Semi)Deviation, Mean-Plus-(Semi)Variance, Entropic Risk, Spectral Risk, Buffered Probability, ...
2. **Distributionally Robust:** Φ-divergence ambiguity including Kullback-Leibler, χ^2, Total Variation, Hellinger, ...
3. **Other Features:** Numerous measures of regret, deviation and error, linear regression tool, stochastic constraints, ...
4. Parallelism and adaptivity through **BatchManager**, **SampleGenerator** and **TrustRegionStep**.

`https://github.com/trilinos/Trilinos`

The Rapid Optimization Library

RAPID OPTIMIZATION LIBRARY

PDE-OPT Application Development Kit



https://github.com/trilinos/Trilinos

The Rapid Optimization Library

RAPID OPTIMIZATION LIBRARY

PDE-OPT Application Development Kit

Physics module	Equations	# Fields	FE space	FE order
Convective heat	Advection-diffusion	1	H(grad)	1 or 2
Radiative heat	Stefan-Boltzmann	1	H(grad)	1 or 2
Semiconductor	Poisson-Boltzmann	1	H(grad)	1 or 2
Structural dynamics	Linear elasticity	d	H(grad)	1 or 2
Incompressible flow	Navier-Stokes	$d+1$	H(grad) + H(div)	1 **and** 2
Thermally convected flow	Navier-Stokes+A.D.	$d+2$	H(grad) + H(div)	1 **and** 2
Superconductivity	Ginzburg-Landau	2	H(grad)	1 or 2
Acoustics	Helmholtz	2	H(grad)	1 or 2

https://github.com/trilinos/Trilinos

Outline

Motivating Applications

General Problem Formulation

Quantifying Risk

Probabilistic Optimization

What if our uncertainty is uncertain?

Computational Solution Methods

Software

References

References – Plus Many More . . .

1. **Ph. Artzner, F. Delbaen, J.-M. Eber & D. Heath,** *Coherent measures of risk.* Math. Finance, 1999.
2. **A. Ben Tal & M. Teboulle**, *Penalty functions and duality in stochastic programming via phi-divergence functionals*, Mathematics of Operations Research, 1987.
3. **A. Ben Tal & M. Teboulle**, *An old-new concept of convex risk measures: The optimized certainty equivalents*, Mathematical Finance, 2007.
4. **D. P. Kouri, M. Heinkenschloss, D. Ridzal, B. G. van Bloemen Waanders**, *A trust-region algorithm with adaptive stochastic collocation for PDE optimization under uncertainty*, SISC, 2013.
5. **D. P. Kouri, M. Heinkenschloss, D. Ridzal, B. G. van Bloemen Waanders**, *Inexact objective function evaluations in a trust-region algorithm for PDE-constrained optimization under uncertainty*, SISC, 2014.
6. **D. P. Kouri & A. Shapiro**, *Optimization of PDEs with Uncertain Inputs*, IMA Volumes in Mathematics and its Applications, 2018.
7. **D. P. Kouri & T. M. Surowiec**, *Existence and optimality conditions for risk-averse PDE-constrained optimization*, SIAM/ASA JUQ (accepted), 2018.
8. **P. A. Krokhmal**, *Higher moment coherent risk measures*, Quantitative Finance, 2007.
9. **S. Kusuoka**, *On law-invariant coherent risk measures*, Advances in Mathematical Economics, 2001.
10. **A. Mafusalov & S. Uryasev**, *Buffered probability of Exceedance: Mathematical Properties and Optimization Algorithms*, UF Technical Report, 2016.
11. **H. Markowitz**, *Portfolio selection*, The Journal of Finance, 1952.
12. **H. Robbins & S. Monro**, *A stochastic approximation method*, Annals of Mathematical Statistics, 1951.
13. **R. T. Rockafellar & S. Uryasev**, *The fundamental risk quadrangle in risk management, optimization, and statistical estimation*, Surveys in Operations Research & Managment Science, 2013.
14. **R. T. Rockafellar & J. O. Royset**, *On buffered failure probability in design and optimization of structures*, Reliability Engineering & System Safety, 2010.
15. **R. T. Rockafellar & J. O. Royset**, *Random variables, monotone relations, and convex analysis*, Mathematical Programming, 2014.
16. **R. T. Rockafellar & R. J.-B. Wets**, *Scenarios and policy aggregation in optimization under uncertainty*, Mathematics of Operations Research, 1991.
17. **W. W. Rogosinski**, *Moments of non-negative mass*, Proceedings of the Royal Society of London: Series A, Mathematical and Physical Sciences, 1958.
18. **A. Shapiro**, *Distributionally robust stochastic programming*, SIAM Journal on Optimization, 2017.
19. **A. Shapiro**, *Monte Carlo sampling methods*, Stochastic Programming, Handbooks in Operations Research and Management Science, 2003.
20. **A. Shapiro, D. Dentcheva & A. Ruszczynski**. *Lectures on Stochastic Programming: Modeling and Theory, Second Edition*, MOS-SIAM Series on Optimization, 2014.